

REDDING

Lolium perenne

- Very early growth start in the spring
- Excellent rust resistance
- Large spring window
- Late heading



Large spring window

With about 64 days from growth start to beginning of heading, REDDING will produce quality forage over a very long period. This makes it a good option for early grazing as the main part of the 64 days comes from the early start compared to varieties heading +/- 3 days after/before REDDING. The early start shows in the high yield of 1st cut too.

Disease tolerant

REDDING has a very high rust tolerance that will preserve forage quality and palatability, especially in the summer and autumn period where this disease can be devastating.

Persistency and quality

Ryegrasses produce high quality forage for demanding animals. REDDING has good persistency and is easy to manage for conservation due to the higher dry matter content than tetraploids.

Technical Specifications

- Perennial ryegrass
- Lating heading diploid perennial ryegrass with early start in spring
- Listed/recommended in EU
DE DK FR

Specification and ratings

Spring window and stems in regrowth

Source : Herbe-book.org 2019

Stems in regrowth: 1 = no stems

Variety	Heading date	Spring start (date)	Spring window (days)	Stems in regrowth
Varieties +/- 3 days	29/5	7/4	59	2,5
Redding	30/5	4/4	64	2,4

Dry matter yield

Source: herbe-book.org 2019

Ton dry matter per hectare

Variety	1st cut	Spring yield	Summer-autumn yield	Total yield
Varieties +/- 3 days	2,0	6,2	2,8	9,1
Redding	2,3	6,2	2,8	9,0

Ratings

Scale 1-9, where 9 = best or most pronounced

